arXiv:hep-th/9904157v1 23 Apr 1999

# The Complete Brane Solution in $D$-dimensional Coupled Gravity System

Bihn Zhou* and Chuan-Jie Zhu†
Institute of Theoretical Physics, Chinese Academy of Sciences, P. O. Box 2735, Beijing 100080, P. R. China

February 1, 2008

**Abstract**

In this letter we present the complete explicit brane solution in $D$-dimensional coupled gravity system.

In a previous paper [1], we studied the brane solutions in $D$-dimensional coupled gravity system to black brane solutions. By using a Poincaré invariant ansatz, we found that we can solve three equations explicitly and there remains only one equation which is a kind of Ricatti equation. Quite recently we have completely solved this last equation. In this letter we report on this complete explicit brane solution. We follow all our previous notations in [1] and refer the reader to this paper for backgrounds and more references.

The equations derived from the coupled $D$-dimensional gravity system are as follows:

$$A'' + d(A')^2 + \tilde{d}\, A'B' + \frac{\tilde{d}+1}{r}\, A' = \frac{\tilde{d}}{2(D-2)}\, S^2, \qquad (1)$$

$$B'' + dA'B' + \frac{d}{r}\, A' + \tilde{d}(B')^2$$

---

*E-mail: zhoub@itp.ac.cn

†E-mail: zhucj@itp.ac.cn

$$+\frac{2\tilde{d}+1}{r}\,B' = -\frac{1}{2}\,\frac{d}{D-2}\,S^2, \tag{2}$$

$$dA'' + (\tilde{d}+1)B'' + d(A')^2 + \frac{\tilde{d}+1}{r}\,B' -dA'B' + \frac{1}{2}\,(\phi')^2 = \frac{1}{2}\,\frac{\tilde{d}}{D-2}\,S^2, \tag{3}$$

$$\phi'' + \Big(dA' + \tilde{d}B' + \frac{\tilde{d}+1}{r}\Big)\phi = -\frac{a}{2}\,S^2, \tag{4}$$

$$(\Lambda'(r)\,e^{\Lambda(r)+a\phi(r)-dA(r)+\tilde{d}B(r)}\,r^{\tilde{d}+1})' = 0, \tag{5}$$

where $S = \Lambda'\,e^{\frac{1}{2}\,a\phi+\Lambda-d\,A}$, $d = p+1 = n-1$ and $\tilde{d} = D-d-2$. These five equations, eqs. (1)-(5), consist of the complete system of equations of motion for four unknown functions: $A(r)$, $B(r)$, $\phi(r)$ and $\Lambda(r)$. We will solve these equations completely.

First it is easy to integrate eq. (5) to get

$$\Lambda'(r)\,e^{\Lambda(r)+a\phi(r)-dA(r)+\tilde{d}B(r)}\,r^{\tilde{d}+1} = C_0, \tag{6}$$

where $C_0$ is a constant of integration. If we know the other three functions $A(r)$, $B(r)$ and $\phi(r)$, this equation can be easily integrated to give $\Lambda(r)$:

$$e^{\Lambda(r)} = C_0\int^r \mathrm{d}r\,\frac{e^{-a\phi(r)+dA(r)-\tilde{d}B(r)}}{r^{\tilde{d}+1}}. \tag{7}$$

By using eq. (6), $S$ can be written as

$$S(r) = C_0\frac{e^{-\frac{a}{2}\phi(r)-\tilde{d}B(r)}}{r^{\tilde{d}+1}}. \tag{8}$$

Now we make a change of functions from $A(r)$, $B(r)$ and $\phi(r)$ to $\xi(r)$, $\eta(r)$ and $Y(r)$:

$$\xi(r) = dA(r) + \tilde{d}B(r), \tag{9}$$

$$\eta(r) = \phi(r) + a(A(r)-B(r)), \tag{10}$$

$$Y(r) = A(r) - B(r), \tag{11}$$

the equations are then changed to

$$\xi'' + (\xi')^2 + \frac{2\tilde{d}+1}{r}\xi' = 0, \tag{12}$$

$$\eta'' + \left(\xi' + \frac{\tilde{d}+1}{r}\right)\eta' - \frac{a}{r}\xi' = 0, \tag{13}$$

$$\begin{aligned} Y'' - \frac{\Delta}{2}(Y')^2 + \left(\frac{\tilde{d}-d}{D-2}\xi' + \frac{\tilde{d}+1}{r} + a\,\eta'\right)Y' \\ -\frac{1}{2}(\eta')^2 - \xi'' + \frac{1}{D-2}(\xi')^2 = 0, \end{aligned} \tag{14}$$

$$Y'' + \left(\xi' + \frac{\tilde{d}+1}{r}\right)Y' - \frac{1}{r}\xi' = \frac{1}{2}S^2, \tag{15}$$

where

$$\Delta = \frac{2d\tilde{d}}{D-2} + a^2. \tag{16}$$

The general solutions for $\xi$ and $\eta$ can be obtained easily from eqs. (12) and (13) and we have

$$\xi = \ln\left|C_1 + C_2 r^{-2\tilde{d}}\right|, \tag{17}$$

$$\eta' = \frac{2C_2 a + C_3 r^{\tilde{d}}}{r(C_2 + C_1 r^{2\tilde{d}})}, \tag{18}$$

where $C_1$, $C_2$ and $C_3$ are constants of integration. Substituting the above expressions into eqs. (14) and (15), we obtain

$$Y'' - \frac{\Delta}{2}(Y')^2 + Q(r)Y' = R(r) \tag{19}$$

and

$$S^2 = \Delta\left(Y' - \frac{2C_2\,\Delta + a\,C_3\,r^{\tilde{d}}}{r\,\Delta\,(C_2 + C_1 r^{2\tilde{d}})}\right)^2 + \frac{K\,r^{2\tilde{d}-2}}{\Delta(C_2 + C_1 r^{2\tilde{d}})^2}, \tag{20}$$

where

$$Q(r) = \frac{\tilde{d}+1}{r} + \frac{2C_2(\Delta - \tilde{d}) + C_3 a r^{\tilde{d}}}{r(C_2 + C_1 r^{2\tilde{d}})}, \tag{21}$$

$$R(r) = \frac{2C_2^2(\Delta - \tilde{d}) + 2C_2C_3 a r^{\tilde{d}} + 2C_1C_2\tilde{d}(2\tilde{d}+1)r^{2\tilde{d}} + \frac{1}{2}C_3^2 r^{2\tilde{d}}}{r^2(C_2 + C_1 r^{2\tilde{d}})^2}, \tag{22}$$

where

$$K = C_3^2(\Delta - a^2) + 8C_1C_2\Delta\tilde{d}(\tilde{d}+1), \tag{23}$$

is a constant.

Now we solve the last equation (19). Setting

$$f = \int dr e^{\Delta Y - \int Q(r) dr}, \tag{24}$$

or,

$$Y = \frac{1}{\Delta}\left(\ln(f') + \int Q(r) dr\right), \tag{25}$$

eq. (19) becomes

$$\left(\frac{f''}{f'}\right)' - \frac{1}{2}\left(\frac{f''}{f'}\right)^2 = \tilde{R}(r), \tag{26}$$

where

$$\begin{aligned} \tilde{R}(r) &= \Delta R(r) - Q'(r) - \frac{1}{2}Q^2(r) \\ &= -\frac{\tilde{d}^2 - 1}{2r^2} + \frac{\tilde{\Lambda} r^{2\tilde{d}-2}}{2(C_2 + C_1 r^{2\tilde{d}})^2} \end{aligned} \tag{27}$$

and

$$\tilde{\Lambda} = K - 4C_1C_2\tilde{d}^2. \tag{28}$$

The left-hand side of eq. (26) is the well-known Schwartz derivative of the function $f$. By exploiting the property of Schwartz derivative [2], a special solution of eq. (26) is given as follows:

$$f_0(r) = \tan\left(k \arctan\sqrt{\frac{C_1}{C_2}}\, r^{\tilde{d}}\right), \tag{29}$$

and the general solution is obtained from this special solution by an arbitrary $SL(2, R)$ transformation:

$$f(r) = \frac{a_0 f_0(r) + b_0}{c_0 f_0(r) + d_0}. \tag{30}$$

Here $k$ is a constant:

$$k = \frac{1}{2\tilde{d}}\sqrt{\frac{K}{C_1C_2}}. \tag{31}$$

The special case considered in [1] corresponds to $k = 1$.

With this general solution in hand one can check that the other equation (20) is also satisfied. The complete solution is rather involved and here we only gave the solution for $f(r) = f_0(r)$:

$$\xi(r) = \ln\left|C_1 + C_2\, r^{-2\tilde{d}}\right|, \tag{32}$$

$$\eta(r) = C_4 - \frac{2\,a}{\tilde{d}}\, \ln\left|C_1 + C_2\, r^{-2\tilde{d}}\right| + \frac{C_3}{\tilde{d}\sqrt{C_1 C_2}}\, \arctan\sqrt{\frac{C_1}{C_2}}\, r^{\tilde{d}}, \tag{33}$$

$$\begin{aligned} Y(r) &= C_5 - \frac{2}{\Delta}\, \ln(\cos(k \arctan\sqrt{\frac{C_1}{C_2}}\, r^{\tilde{d}})) \\ &\quad -\frac{1}{\tilde{d}}\, \ln\left|C_1 + C_2\, r^{-2\tilde{d}}\right| + \frac{a\, C_3}{\tilde{d}\Delta\sqrt{C_1 C_2}}\, \arctan\sqrt{\frac{C_1}{C_2}}\, r^{\tilde{d}}, \end{aligned} \tag{34}$$

and

$$\begin{aligned} A(r) &= \frac{\tilde{d}\, C_5}{D-2} - \frac{2\tilde{d}}{(D-2)\Delta}\, \ln(\cos(k \arctan\sqrt{\frac{C_1}{C_2}}\, r^{\tilde{d}})) \\ &\quad +\frac{a\, C_3}{(D-2)\Delta\sqrt{C_1\, C_2}}\, \arctan\sqrt{\frac{C_1}{C_2}}\, r^{\tilde{d}}, \end{aligned} \tag{35}$$

$$\begin{aligned} B(r) &= -\frac{d\, C_5}{D-2} + \frac{2\, d}{(D-2)\Delta}\, \ln(\cos(k \arctan\sqrt{\frac{C_1}{C_2}}\, r^{\tilde{d}})) \\ &\quad +\frac{1}{\tilde{d}}\, \ln\left|C_1 + C_2\, r^{-2\tilde{d}}\right| - \frac{a\, d\, C_3}{\tilde{d}\,(D-2)\Delta\sqrt{C_1 C_2}}\, \arctan\sqrt{\frac{C_1}{C_2}}\, r^{\tilde{d}}, \end{aligned} \tag{36}$$

$$\begin{aligned} \phi(r) &= C_4 + \frac{2\, a}{\Delta}\, \ln(\cos(k \arctan\sqrt{\frac{C_1}{C_2}}\, r^{\tilde{d}})) \\ &\quad +\frac{2\, d\, C_3}{(D-2)\Delta\sqrt{C_1 C_2}} \arctan\sqrt{\frac{C_1}{C_2}}\, r^{\tilde{d}}, \end{aligned} \tag{37}$$

$$e^{\Lambda(r)} = C_0\, e^{-aC_4+(\Delta-a^2)C_5}\, \left[\frac{2}{\sqrt{K}}\, \tan(k \arctan\sqrt{\frac{C_1}{C_2}}\, r^{\tilde{d}}) - C_6\right], \tag{38}$$

with

$$C_0 = e^{\frac{a}{2}\, C_4 - \frac{1}{2}\,(\Delta-a^2)\, C_5}\, \sqrt{\frac{K}{\Delta}}. \tag{39}$$

More details and the extension of the complete explicit solution to black branes will be presented elsewhere [2, 3].

## Acknowledgments

We would like to thank Han-Ying Guo, Yi-hong Gao, Ke Wu, Ming Yu, Zhu-jun Zheng and Zhong-Yuan Zhu for discussions. This work is supported in part by funds from Chinese National Science Foundation and Pandeng Project.

## References

[1] B. Zhou and C. -J. Zhu, *A Study of Brane Solutions in D-Dimensional Coupled Gravity System*, preprint hep-th/9903118.

[2] B. Zhou and C. -J. Zhu, *The Complete Black Brane Solutions in D-Dimensional Coupled Gravity System*, to appear.

[3] B. Zhou, ITP Doctoral Thesis, 1999.